

Prophylactic Antibiotic Regimen Selection for Surgery Specifications Manual 4.0 01/01/2012 through 06/30/2012 Discharges		Prophylactic Antibiotic Regimen Selection for Surgery Specifications Manual 4.0 01/01/2012 through 06/30/2012 Discharges	
Surgical Procedure	Approved Antibiotics	Surgical Procedure	Approved Antibiotics
<b>CABG, Other Cardiac or Vascular</b>	Cefazolin, Cefuroxime, Table 3.1 or Vancomycin <sup>1</sup> Table 3.8  <b>If β-lactam allergy:</b> Vancomycin <sup>2</sup> Table 3.8 or Clindamycin <sup>2</sup> Table 3.9	<b>Hysterectomy</b>	Cefotetan, Cefazolin, Cefoxitin, Cefuroxime, or Ampicillin/Sulbactam Table 3.7 <b>If β-lactam allergy:</b> Clindamycin Table 3.9 + Aminoglycoside Table 2.11, or Clindamycin Table 3.9 + Quinolone Table 3.12, or Clindamycin Table 3.9 + Aztreonam Table 2.7 <b>OR</b> Metronidazole Table 3.6a + Aminoglycoside Table 2.11, or Metronidazole Table 3.6a + Quinolone Table 3.12
<b>Hip/Knee Arthroplasty</b>	Cefazolin or Cefuroxime Table 3.2 or Vancomycin <sup>1</sup> Table 3.8  <b>If β-lactam allergy:</b> Vancomycin <sup>2</sup> Table 3.8 or Clindamycin <sup>2</sup> Table 3.9	<b>Principal Procedure Code of Hysterectomy with an Other Procedure Code of Colon Surgery found in Appendix A, Table 5.03</b>	Cefotetan, Cefazolin, Cefoxitin, Cefuroxime, or Ampicillin/Sulbactam Table 3.7 <b>OR</b> Ertapenem <sup>3</sup> Table 3.6b <b>If β-lactam allergy:</b> Clindamycin Table 3.9 + Aminoglycoside Table 2.11, or Clindamycin Table 3.9 + Quinolone Table 3.12, or Clindamycin Table 3.9 + Aztreonam Table 2.7 <b>OR</b> Metronidazole Table 3.6a + Aminoglycoside Table 2.11, or Metronidazole Table 3.6a + Quinolone Table 3.12
<b>Colon</b>	Cefotetan, Cefoxitin, Ampicillin/Sulbactam Table 3.5, or Ertapenem <sup>3</sup> Table 3.6b <b>OR</b> Cefazolin or Cefuroxime Table 3.2 + Metronidazole Table 3.6a <b>If β-lactam allergy:</b> Clindamycin Table 3.9 + Aminoglycoside Table 2.11, or Clindamycin Table 3.9 + Quinolone Table 3.12, or Clindamycin Table 3.9 + Aztreonam Table 2.7 <b>OR</b> Metronidazole Table 3.6a with Aminoglycoside Table 2.11, or Metronidazole Table 3.6a + Quinolone Table 3.12	<b>Special Considerations:</b> <ol style="list-style-type: none"> <li><b>Vancomycin is acceptable with a physician/APN/PA/pharmacist documented justification for its use (see data element Vancomycin).</b></li> <li><b>For cardiac, orthopedic, and vascular surgery, if the patient is allergic to beta-lactam antibiotics, Vancomycin or Clindamycin are acceptable substitutes.</b></li> <li><b>A single dose of Ertapenem is recommended for colon procedures.</b></li> </ol>	



This material, developed by FMQAI, the Medicare Quality Improvement Organization (QIO) for Florida, and Health Services Advisory Group of California, Inc., the Medicare QIO for California, under contract with the Centers for Medicare & Medicaid Services (CMS), an agency of the U.S. Department of Health and Human Services. The contents presented do not necessarily reflect CMS policy.  
FL-10SOW-2011FLC706-11-12539, CA-10SOW-7.4-111011-02

Prophylactic Antibiotic Regimen Selection for Surgery Specifications Manual 4.0 01/01/2012 through 06/30/2012 Discharges		Prophylactic Antibiotic Regimen Selection for Surgery Specifications Manual 4.0 01/01/2012 through 06/30/2012 Discharges	
Surgical Procedure	Approved Antibiotics	Surgical Procedure	Approved Antibiotics
<b>CABG, Other Cardiac or Vascular</b>	Cefazolin, Cefuroxime, Table 3.1 or Vancomycin <sup>1</sup> Table 3.8  <b>If β-lactam allergy:</b> Vancomycin <sup>2</sup> Table 3.8 or Clindamycin <sup>2</sup> Table 3.9	<b>Hysterectomy</b>	Cefotetan, Cefazolin, Cefoxitin, Cefuroxime, or Ampicillin/Sulbactam Table 3.7 <b>If β-lactam allergy:</b> Clindamycin Table 3.9 + Aminoglycoside Table 2.11, or Clindamycin Table 3.9 + Quinolone Table 3.12, or Clindamycin Table 3.9 + Aztreonam Table 2.7 <b>OR</b> Metronidazole Table 3.6a + Aminoglycoside Table 2.11, or Metronidazole Table 3.6a + Quinolone Table 3.12
<b>Hip/Knee Arthroplasty</b>	Cefazolin or Cefuroxime Table 3.2 or Vancomycin <sup>1</sup> Table 3.8  <b>If β-lactam allergy:</b> Vancomycin <sup>2</sup> Table 3.8 or Clindamycin <sup>2</sup> Table 3.9	<b>Principal Procedure Code of Hysterectomy with an Other Procedure Code of Colon Surgery found in Appendix A, Table 5.03</b>	Cefotetan, Cefazolin, Cefoxitin, Cefuroxime, or Ampicillin/Sulbactam Table 3.7 <b>OR</b> Ertapenem <sup>3</sup> Table 3.6b <b>If β-lactam allergy:</b> Clindamycin Table 3.9 + Aminoglycoside Table 2.11, or Clindamycin Table 3.9 + Quinolone Table 3.12, or Clindamycin Table 3.9 + Aztreonam Table 2.7 <b>OR</b> Metronidazole Table 3.6a + Aminoglycoside Table 2.11, or Metronidazole Table 3.6a + Quinolone Table 3.12
<b>Colon</b>	Cefotetan, Cefoxitin, Ampicillin/Sulbactam Table 3.5, or Ertapenem <sup>3</sup> Table 3.6b <b>OR</b> Cefazolin or Cefuroxime Table 3.2 + Metronidazole Table 3.6a <b>If β-lactam allergy:</b> Clindamycin Table 3.9 + Aminoglycoside Table 2.11, or Clindamycin Table 3.9 + Quinolone Table 3.12, or Clindamycin Table 3.9 + Aztreonam Table 2.7 <b>OR</b> Metronidazole Table 3.6a with Aminoglycoside Table 2.11, or Metronidazole Table 3.6a + Quinolone Table 3.12	<b>Special Considerations:</b> <ol style="list-style-type: none"> <li><b>Vancomycin is acceptable with a physician/APN/PA/pharmacist documented justification for its use (see data element Vancomycin).</b></li> <li><b>For cardiac, orthopedic, and vascular surgery, if the patient is allergic to beta-lactam antibiotics, Vancomycin or Clindamycin are acceptable substitutes.</b></li> <li><b>A single dose of Ertapenem is recommended for colon procedures.</b></li> </ol>	



This material, developed by FMQAI, the Medicare Quality Improvement Organization (QIO) for Florida, and Health Services Advisory Group of California, Inc., the Medicare QIO for California, under contract with the Centers for Medicare & Medicaid Services (CMS), an agency of the U.S. Department of Health and Human Services. The contents presented do not necessarily reflect CMS policy.  
FL-10SOW-2011FLC706-11-12539, CA-10SOW-7.4-111011-02